

AASC™

Agitation in Alzheimer's Screener for Caregivers

A person with Alzheimer's dementia may become easily upset, frustrated, or angry. These feelings may be associated with agitation, a common symptom of Alzheimer's.

The following questions ask about behaviors associated with agitation that you may have observed in the individual for whom you provide care.

Your answers to these questions can help you talk with your person's healthcare professional. Please select one response for each question.

Patient Name _____ Date _____

1. Are you noticing any of the following that represent a change from the individual's usual or past behavior?

	Yes	No
a. Repeating motions or behaviors (e.g., rocking, raising fist, pointing finger)	<input type="checkbox"/>	<input type="checkbox"/>
b. Pacing or restlessness (cannot be still)	<input type="checkbox"/>	<input type="checkbox"/>
c. Cursing/using profanity or lashing out verbally	<input type="checkbox"/>	<input type="checkbox"/>
d. Raising voice or yelling or screaming	<input type="checkbox"/>	<input type="checkbox"/>
e. Resisting assistance or care	<input type="checkbox"/>	<input type="checkbox"/>
f. Throwing or hitting or breaking things	<input type="checkbox"/>	<input type="checkbox"/>
g. Trying to hurt self or others (e.g., grabbing, kicking, hitting, biting)	<input type="checkbox"/>	<input type="checkbox"/>

2. Do any of these behaviors negatively affect the individual's relationships, activities, or willingness to receive care?

Yes	No
<input type="checkbox"/>	<input type="checkbox"/>

Disclaimer: This screener was developed by Otsuka Global Medical Affairs along with a panel of multidisciplinary experts and was funded by Otsuka Pharmaceutical Development & Commercialization, Inc. The AASC™ is a screening tool and is not a diagnostic tool.

PsychU is supported by Otsuka Pharmaceutical Development & Commercialization, Inc. (OPDC), Otsuka America Pharmaceutical, Inc. (OAPI), and Lundbeck, LLC – committed supporters of the mental health treatment community. The opinions expressed by PsychU's contributors are their own and are not endorsed or recommended by PsychU or its sponsors. The information provided through PsychU is intended for the educational benefit of mental health care professionals and others who support mental health care. It is not intended as, nor is it a substitute for, medical care, advice, or professional diagnosis. Health care professionals should use their independent medical judgement when reviewing PsychU's educational resources. Users seeking medical advice should consult with a health care professional. No CME or CEU credits are available through any of the resources provided by PsychU. Some of the contributors may be paid consultants for OPDC, OAPI, and / or Lundbeck, LLC.

Agitation criteria based on the International Psychogeriatric Association (IPA) definition (Sano et al., 2023).

References: 1. Clevenger C, et al. One Minute to Recognition: The Agitation in Alzheimer's Screener for Caregivers (AASC™). The Gerontological Society of America Annual Scientific Meeting; November 8-12, 2023; Tampa, FL. 2. Sano M, Cummings J, Auer S, et al. *Int Psychogeriatr.* 2023;1-13.

© 2024 Otsuka Pharmaceutical Development & Commercialization, Inc. All rights reserved.